

Regular vertex operator subalgebras and compressions of intertwining operators

BIN GUI

Abstract

Let V be a vertex operator subalgebra of U . Assume that U , V , and its commutant V^c in U are CFT-type, self-dual, and regular VOAs. Assume also that the double commutant V^{cc} equals V . We prove that any intertwining operator of V is a compression of intertwining operators of U .

0 Introduction

In [KM15], Krauel-Miyamoto showed that if V is a vertex operator subalgebra of U , if U , V , and the commutant V^c are CFT-type, self-dual, and regular VOAs, and if $V^{cc} = V$, then any irreducible V -module appears in some irreducible U -module. For example, by [Ara15a, Ara15b, ACL19], these assumptions are satisfied when we take $V \subset U$ to be $L_{k+1}(\mathfrak{g}) \subset L_k(\mathfrak{g}) \otimes L_1(\mathfrak{g})$, where k is a positive integer, \mathfrak{g} is a finite dimensional complex simple Lie algebra of type ADE , and $L_k(\mathfrak{g})$ is the corresponding (unitary) affine VOA. In this case, V^c is a discrete series principle W -algebra $\mathcal{W}_l(\mathfrak{g})$.

The above results have important applications to the unitarity problems in VOAs: For example, we can conclude that any irreducible $\mathcal{W}_l(\mathfrak{g})$ -module is unitarizable since this is true for any unitary affine VOA. Moreover, using these results (together with the techniques developed in [Ten17, Ten19a, Ten19b]), Tener showed in [Ten19c] that the modular tensor categories associated to all unitary affine VOAs and type AE discrete series W -algebras are unitary, and solved a longstanding problem in subfactor theory and algebraic quantum field theory: that the conformal nets associated to unitary affine VOAs and type ADE discrete series W -algebras are completely rational.

In this paper, we generalize the result of [KM15] to intertwining operators: We show that any intertwining operator of V is a compression of intertwining operators of U (theorem 4.4). To be more precise, let W_I, W_J, W_K be (ordinary) U -modules, which can also be regarded as weak V -modules. Suppose that \mathcal{Y}^U is a type $\binom{W_K}{W_I W_J}$ intertwining operator of U , and W_i, W_j, W_k are graded irreducible V -submodules of W_I, W_J, W_K respectively, then one can find $\lambda \in \mathbb{Q}$ such that z^λ times the restriction of \mathcal{Y}^U to W_i, W_j, W_k is an intertwining operator \mathcal{Y} of V . (Note that without the factor z^λ , the restriction itself may not satisfy the L_{-1} -derivative property.) We then say that \mathcal{Y} is a **compression** of \mathcal{Y}^U . (See definition 3.4 for more details.) Our main result of this article is that any intertwining operator of V can be written as a (finite) sum of those that are compressions of intertwining operators of U .

Our result can be applied to prove many important functional analytic properties for intertwining operators. One such property is the (polynomial) energy bounds condition [CKLW18, Gui19a, Gui19b], which says roughly that the smeared intertwining operators are bounded by L_0^n for some $n \geq 0$. Proving energy bounds condition for intertwining operators is a key step in relating the tensor structures of VOA modules and the corresponding conformal net modules; see [Was98, TL04, Gui18, Gui20]. On the other hand, one may deduce the energy bounds condition of the compressed intertwining operator \mathcal{Y} from that of \mathcal{Y}^U . Since, by our main result, any intertwining operator of $L_k(\mathfrak{g})$ or $\mathcal{W}_l(\mathfrak{g})$ (when \mathfrak{g} is of type *ADE*) is a compression of tensor products of intertwining operators of $L_1(\mathfrak{g})$, and since the latter were proved in [TL04] to be energy bounded, we can conclude that all intertwining operators of $L_k(\mathfrak{g})$ or $\mathcal{W}_l(\mathfrak{g})$ are energy bounded. This result will be used in [Gui20] to show that if V is $L_k(\mathfrak{g})$ or $\mathcal{W}_l(\mathfrak{g})$, and if \mathcal{A} is the corresponding conformal net, then the tensor and braid structures of the representation categories of V and \mathcal{A} are compatible.

Acknowledgment

The author would like to thank James Tener for helpful discussions on the topic of this paper.

1 Intertwining operators and tensor categories

Let $(V, Y, \mathbf{1}, \nu)$ be a vertex operator algebra (VOA) where $\mathbf{1}$ is the vacuum vector and ν is the conformal vector. For any $v \in V$, write $Y(v, z) = \sum_{n \in \mathbb{Z}} Y(v)_n z^{-n-1}$ where $Y(v)_n \in \text{End}(V)$. Then $L_n := Y(\nu)_{n+1}$ satisfy the Virasoro relation

$$[L_n, L_m] = (n - m)L_{n+m} + \delta_{n,-m} \frac{n^3 - n}{12} c,$$

where c is the central charge of V . We shall always assume that V is CFT-type, namely, V has L_0 -grading $V = \bigoplus_{n \in \mathbb{N}} V(n)$ and $V(0) = \mathbb{C}\mathbf{1}$. We also assume that V is self-dual and regular (equivalently, self-dual, rational and C_2 -cofinite [ABD04]). Note that the self-dual condition is equivalent to the existence of a non-degenerate invariant bilinear form. As a consequence of CFT-type and being self-dual, V is simple. (See, for example, [CKLW18] proposition 6.4-(iv).) Moreover, any (ordinary) V -module is semisimple, and the category of V -modules is a rigid modular tensor category [Hua08].

We write V -modules as W_i, W_j, W_k, \dots whose vertex operators are denoted by Y_i, Y_j, Y_k, \dots respectively. V itself as a V -module (the vacuum module) will also be written as W_0 . We write $Y_i(v, z) = \sum_{n \in \mathbb{Z}} Y_i(v)_n z^{-n-1}$ where each $Y_i(v)_n \in \text{End}(W_i)$. Again, any V -module has L_0 -grading $W_i = \bigoplus_{n \in \mathbb{C}} W_i(n)$. Recall that a homogeneous vector of W_i is, by definition, an eigenvector of L_0 . In the case that W_i is irreducible (i.e. simple), we furthermore have $W_i = \bigoplus_{n \in \mathbb{N} + \alpha} W_i(n)$ for some $\alpha \in \mathbb{C}$. (Indeed, $\alpha \in \mathbb{Q}$ by [AM88, DLM00].) We let P_n denote the projection of W_i onto $W_i(n)$. We also let

$$W_i(\leq n) = \bigoplus_{\text{Re}(m) \leq n} W_i(m),$$

and let $P_{\leq n}$ be the projection of W_i onto $W_i(\leq n)$.

Let $W_{\bar{i}}$ denote the contragredient module of W_i . Recall that as a vector space, $W_{\bar{i}} = \bigoplus_{n \in \mathbb{C}} W_i(n)^*$. (See [FHL93] for more details.) The evaluation between $w' \in W_{\bar{i}}$ and $w \in W_i$ is written as $\langle w, w' \rangle$ or $\langle w', w \rangle$. (The same notation will be used if one of w, w' is in the algebraic completion.) Since V is self-dual, we identify the vacuum module $V = W_0$ and its contragredient module $W_{\bar{0}}$. $W_{\bar{i}}$ is identified with W_i in an obvious way.

Recall that if W_i, W_j, W_k are V -modules, an intertwining operator \mathcal{Y} of type $\binom{W_k}{W_i W_j}$ (or $\binom{k}{i j}$ for short) is a linear map

$$\begin{aligned} W_i &\rightarrow \text{End}(W_j, W_k)\{z\} \\ w_i &\mapsto \mathcal{Y}(w^{(i)}, z) = \sum_{n \in \mathbb{C}} \mathcal{Y}(w^{(i)})_n z^{-n-1} \end{aligned}$$

where the sum above is the formal sum, each $\mathcal{Y}(w^{(i)})_n$ is in $\text{End}(W_j, W_k)$, and the following conditions are satisfied:

(a) (Lower truncation) For any $w^{(j)} \in W_j$, $\mathcal{Y}(w^{(i)})_n w^{(j)} = 0$ when $\text{Re}(n)$ is sufficiently large.

(b) (Jacobi identity) For any $u \in V, w^{(i)} \in W_i, m, n \in \mathbb{Z}, s \in \mathbb{C}$, we have

$$\begin{aligned} &\sum_{l \in \mathbb{N}} \binom{m}{l} \mathcal{Y}(Y_i(u)_{n+l} w^{(i)})_{m+s-l} \\ &= \sum_{l \in \mathbb{N}} (-1)^l \binom{n}{l} Y_k(u)_{m+n-l} \mathcal{Y}(w^{(i)})_{s+l} - \sum_{l \in \mathbb{N}} (-1)^{l+n} \binom{n}{l} \mathcal{Y}(w^{(i)})_{n+s-l} Y_j(u)_{m+l}. \end{aligned} \quad (1.1)$$

(c) (L_{-1} -derivative) $\frac{d}{dz} \mathcal{Y}(w^{(i)}, z) = \mathcal{Y}(L_{-1} w^{(i)}, z)$ for any $w^{(i)} \in W_i$.

We say that W_i, W_j, W_k are respectively the **charge space**, the **source space**, and the **target space** of \mathcal{Y} . If W_i, W_j, W_k are all irreducible, we say that \mathcal{Y} is an **irreducible intertwining operator**.

Recall that given $\mathcal{Y} \in \mathcal{V} \binom{k}{i j}$, one can define $\mathbb{B}\mathcal{Y} \in \mathcal{V} \binom{k}{j i}$ and $\mathbb{C}\mathcal{Y} \in \mathcal{V} \binom{\bar{j}}{i \bar{k}}$, called the (positively) **braided** intertwining operator and the **contragredient** intertwining operator of \mathcal{Y} , by choosing any $w^{(i)} \in W_i, w^{(j)} \in W_j, w^{(\bar{k})} \in W_{\bar{k}}$ and setting

$$\begin{aligned} \mathbb{B}\mathcal{Y}(w^{(j)}, z) w^{(i)} &= e^{zL_{-1}} \mathcal{Y}(w^{(i)}, e^{i\pi} z) w^{(j)}, \\ \langle \mathbb{C}\mathcal{Y}(w^{(i)}, z) w^{(\bar{k})}, w^{(j)} \rangle &= \langle w^{(\bar{k})}, \mathcal{Y}(e^{zL_1}(e^{i\pi} z^{-2})^{L_0} w^{(i)}, z^{-1}) w^{(j)} \rangle. \end{aligned}$$

In particular, $Y_{\bar{i}} \in \mathcal{V} \binom{\bar{i}}{0 \bar{i}}$ (the vertex operator for $W_{\bar{i}}$) is contragredient to Y_i . See [FHL93] for details.

We refer the reader to [HL13] and the references therein for the definition and basic properties of the tensor category $\text{Rep}(V)$ of V -modules. (See also [Gui19a] for a sketch of the Huang-Lepowsky tensor product theory.) Roughly speaking, $\text{Rep}(V)$ is defined such that the fusion rules are exactly the dimensions of the spaces of intertwining operators, and the R - and F -matrices are described by the braid and the fusion relations of intertwining operators.

To be more precise, the tensor functor (fusion product) \boxtimes is defined in such a way that there is a functorial isomorphism

$$\text{Hom}_V(W_i \boxtimes W_j, W_k) \xrightarrow{\cong} \mathcal{V} \binom{k}{i j}, \quad \alpha \mapsto \mathcal{Y}_\alpha \quad (1.2)$$

for any V -modules W_i, W_j, W_k . By saying this map is functorial, we mean that if $F \in \text{Hom}_V(W_{i'}, W_i)$, $G \in \text{Hom}_V(W_{j'}, W_j)$, and $H \in \text{Hom}_V(W_k, W_{k'})$, then for any $w^{(i')} \in W_{i'}$ and $w^{(j')} \in W_{j'}$,

$$\mathcal{Y}_{H\alpha(F\otimes G)}(w^{(i')}, z)w^{(j')} = H\mathcal{Y}_\alpha(Fw^{(i')}, z)Gw^{(j')}.$$

In particular, we denote by $\mathcal{L}_{i,j} \in \mathcal{V}\binom{i \boxtimes j}{i \ j} = \mathcal{V}\binom{W_i \boxtimes W_j}{W_i \ W_j}$ the value of the identity element $\mathbf{1}_{i \boxtimes j} \in \text{Hom}_V(W_i \boxtimes W_j, W_i \boxtimes W_j)$ under (1.2), i.e.,

$$\mathcal{L}_{i,j} = \mathcal{Y}_{\mathbf{1}_{i \boxtimes j}}.$$

Here we adopt the notation

$$W_{i \boxtimes j} = W_i \boxtimes W_j.$$

Then for any $\alpha \in \text{Hom}_V(W_i \boxtimes W_j, W_k)$, by the functoriality of (1.2) and that $\alpha = \alpha \cdot \mathbf{1}_{W_i \boxtimes W_j}$, it is clear that

$$\mathcal{Y}_\alpha = \alpha \mathcal{L}_{i,j}.$$

(In the papers of Huang-Lepowsky, $\mathcal{L}_{i,j}(w^{(i)}, z)w^{(j)}$ is written as $w^{(i)} \boxtimes_{P(z)} w^{(j)}$, regarded as a fusion product of the vectors $w^{(i)}, w^{(j)}$.)

Note that $Y_i \in \mathcal{V}\binom{i}{0 \ i}$. The left unitor $W_0 \boxtimes W_i \xrightarrow{\cong} W_i$ is defined such that it is sent by (1.2) to the element Y_i . The right unitor

$$\kappa(i) : W_i \boxtimes W_0 \xrightarrow{\cong} W_i$$

is defined such that

$$\mathcal{Y}_{\kappa(i)} = \mathbb{B}Y_i.$$

We call $\mathcal{Y}_{\kappa(i)} \in \mathcal{V}\binom{i}{i \ 0}$ the **creation operator** of W_i .

The braid isomorphism $\mathbb{B} = \mathbb{B}_{i,j} : W_i \boxtimes W_j \xrightarrow{\cong} W_j \boxtimes W_i$ is defined such that

$$\mathcal{Y}_{\mathbb{B}_{i,j}} = \mathbb{B}\mathcal{L}_{i,j}.$$

We will not use braiding in this article. To describe the associativity isomorphisms, we first notice:

Proposition 1.1. *Choose any $z \in \mathbb{C}^\times = \mathbb{C} - \{0\}$. Then for any $n \in \mathbb{C}$,*

$$\text{Span}_{w^{(i)} \in W_i, w^{(j)} \in W_j} P_{\leq n} \cdot \mathcal{L}_{i,j}(w^{(i)}, z)w^{(j)} = (W_i \boxtimes W_j)(\leq n). \quad (1.3)$$

This proposition was proved in [Gui19a] section A.2.¹ The main idea of the proof is as follows. Set $W_k = W_i \boxtimes W_j$. Then it is equivalent to proving that for any $w^{(\bar{k})} \in W_{\bar{k}}$, if

$$\langle w^{(\bar{k})}, \mathcal{L}_{i,j}(w^{(i)}, z)w^{(j)} \rangle = 0$$

¹Proposition 1.1 is similar to but slightly stronger than [Hua95] Lemma 14.9. That lemma says that (1.3) holds with $P_{\leq s}$ replaced by P_s and $(W_i \boxtimes W_j)(\leq s)$ by $(W_i \boxtimes W_j)(s)$. Huang's result is enough for applications in our paper.

for any $w^{(i)} \in W_i$ and $w^{(j)} \in W_j$, then $w^{(\bar{k})} = 0$. To see this, let $\mathcal{W} \subset W_{\bar{k}}$ be the subspace of all $w^{(\bar{k})}$ satisfying the above identity. Using the Jacobi identity for intertwining operators, it is easy to see that \mathcal{W} is V -invariant, i.e. it is a V -submodule of $W_{\bar{k}}$. Assume \mathcal{W} is non-trivial. Choose an irreducible module W_l such that \mathcal{W} has an irreducible submodule isomorphic to W_l . Then there is a non-zero $T \in \text{Hom}_V(W_k, W_l)$ whose transpose $T^t \in \text{Hom}_V(W_l, W_{\bar{k}})$ maps W_l into \mathcal{W} . Using the definition of \mathcal{W} and the fact that $T^t w^{(\bar{l})} \in \mathcal{W}$ for each $w^{(\bar{l})} \in W_{\bar{l}}$, it is easy to see that $T\mathcal{L}_{i,j} = 0$. Recall $W_k = W_i \boxtimes W_j$. So we can write $T\mathcal{L}_{i,j} = \mathcal{Y}_T$. Therefore $T = 0$, which gives a contradiction.

Corollary 1.2. *Let W_i, W_j, W_s be V -modules, and assume that W_s is isomorphic to an irreducible submodule of $W_i \boxtimes W_j$. If $\Xi_{i,j}^s$ is a basis of $\text{Hom}_V(W_i \boxtimes W_j, W_s)$, $\alpha \in \Xi_{i,j}^s$, then there exist homogeneous vectors $w_1^{(i)}, \dots, w_m^{(i)} \in W_i$, $w_1^{(j)}, \dots, w_m^{(j)} \in W_j$, $w^{(\bar{s})} \in W_{\bar{s}}$, and constants $\lambda_1, \dots, \lambda_m \in \mathbb{Q}$, such that for any $\beta \in \Xi_{i,j}^s$, the expression*

$$\sum_{l=1}^m z^{\lambda_l} \langle \mathcal{Y}_\beta(w_l^{(i)}, z) w_l^{(j)}, w^{(\bar{s})} \rangle \quad (1.4)$$

is a constant (where z is a complex variable), and it is non-zero if and only if $\beta = \alpha$.

Proof. Choose $n \in \mathbb{C}$ such that $W_s(n)$ is non-trivial. Choose for each $\alpha \in \Xi_{i,j}^s$ a morphism $\check{\alpha} \in \text{Hom}_V(W_s, W_i \boxtimes W_j)$ such that $\check{\alpha}\beta = \delta_{\alpha,\beta} \mathbf{1}_s$ for any $\alpha, \beta \in \Xi_{i,j}^s$. Now we fix $\alpha \in \Xi_{i,j}^s$, and choose a non-zero vector $w^{(\bar{s})} \in W_s(n)$. By proposition 1.1, there exist homogeneous vectors $w_1^{(i)}, \dots, w_m^{(i)} \in W_i$ and $w_1^{(j)}, \dots, w_m^{(j)} \in W_j$ such that

$$\sum_{l=1}^m P_n \mathcal{L}_{i,j}(w_l^{(i)}, 1) w_l^{(j)} = \check{\alpha} w^{(\bar{s})}.$$

Choose $w^{(\bar{s})} \in W_{\bar{s}}(n)$ such that $\langle w^{(s)}, w^{(\bar{s})} \rangle \neq 0$. (Recall that $W_{\bar{s}}(n)$ is the dual vector space of $W_s(n)$.) Then it is easy to check that the constant

$$\sum_{l=1}^m \langle \mathcal{Y}_\beta(w_l^{(i)}, 1) w_l^{(j)}, w^{(\bar{s})} \rangle \quad (1.5)$$

is non-zero (in which case equals $\langle w^{(s)}, w^{(\bar{s})} \rangle$) if and only if $\beta = \alpha$. Now let

$$\lambda_l = \text{wt}(w_l^{(s)}) - \text{wt}(w_l^{(i)}) - \text{wt}(w_l^{(j)}),$$

where the three terms on the right hand side are the weights (i.e. the L_0 -eigenvalues) of the corresponding homogeneous vectors. Then (1.5) equals (1.4). This proves the claim of this corollary. \square

For any irreducible equivalence class of V -modules we choose a representing element and let them form a (finite) set \mathcal{E} . We assume $W_0 \in \mathcal{E}$. Choose V -modules W_i, W_j, W_k, W_l . If $W_s \in \mathcal{E}$, we write $s \in \mathcal{E}$ for short. For each $r \in \mathcal{E}$, we choose bases $\Xi_{j,k}^r$ of $\text{Hom}_V(W_j \boxtimes W_k, W_r)$ and $\Xi_{i,r}^l$ of $\text{Hom}_V(W_i \boxtimes W_r, W_l)$. Choose $z, \zeta \in \mathbb{C}^\times$ with $0 < |\zeta| < |z|$. Then for any $w^{(i)} \in W_i$, $w^{(j)} \in W_j$, and any $\alpha \in \Xi_{i,r}^l$, $\beta \in \Xi_{j,k}^r$ the product

$$\mathcal{Y}_\alpha(w^{(i)}, z) \mathcal{Y}_\beta(w^{(j)}, \zeta) \quad (1.6)$$

converges absolutely [Hua05], in the sense that for any $w^{(k)} \in W_k, w^{(\bar{l})} \in W_{\bar{l}}$,

$$\sum_{n \in \mathbb{C}} |\langle \mathcal{Y}_\alpha(w^{(i)}, z) P_n \mathcal{Y}_\beta(w^{(j)}, \zeta) w^{(k)}, w^{(\bar{l})} \rangle| < +\infty.$$

Moreover, consider the expression

$$\langle \mathcal{Y}_\alpha(w^{(i)}, z) \mathcal{Y}_\beta(w^{(j)}, \zeta) w^{(k)}, w^{(\bar{l})} \rangle \quad (1.7)$$

as an element of $(W_i \otimes W_j \otimes W_k \otimes W_{\bar{l}})^*$. (Note that this element depends also on the arguments $\arg z$ and $\arg \zeta$.) By linearity, we have a linear map

$$\Psi_{z,\zeta} : \bigoplus_{r \in \mathcal{E}} \text{Hom}_V(W_i \boxtimes W_r, W_l) \otimes \text{Hom}_V(W_j \boxtimes W_k, W_r) \rightarrow (W_i \otimes W_j \otimes W_k \otimes W_{\bar{l}})^* \quad (1.8)$$

sending $\alpha \otimes \beta$ to the linear functional defined by (1.7). This map is well-known to be injective. Indeed, choose any \mathfrak{X} in the domain of $\Psi_{z,\zeta}$. If $\Psi_{z,\zeta}(\mathfrak{X})$ equals 0 for one pair (z, ζ) , then, by the existence of differential equations as in [Hua05], $\Psi_{z,\zeta}(\mathfrak{X})$ equals 0 for all z, ζ satisfying $0 < |\zeta| < |z|$. (See [Gui19a] the paragraphs after theorem 2.4 for a detailed explanation.) Then, using proposition 1.1, it is not hard to show $\mathfrak{X} = 0$. (See for instance [Gui19a] proposition 2.3 whose proof is in section A.2.)

Similarly, when $0 < |z - \zeta| < |\zeta|$, one can define an injective linear map

$$\Phi_{z,\zeta} : \bigoplus_{s \in \mathcal{E}} \text{Hom}_V(W_s \boxtimes W_k, W_l) \otimes \text{Hom}_V(W_i \boxtimes W_j, W_s) \rightarrow (W_i \otimes W_j \otimes W_k \otimes W_{\bar{l}})^* \quad (1.9)$$

sending each $\gamma \otimes \delta$ to the linear functional determined by the following iterate of intertwining operators:

$$\langle \mathcal{Y}_\gamma(\mathcal{Y}_\delta(w^{(i)}, z - \zeta) w^{(j)}, \zeta) w^{(k)}, w^{(\bar{l})} \rangle. \quad (1.10)$$

Again, this map depends on the choice of arguments: $\arg(z - \zeta)$ and $\arg(\zeta)$, and the above expression converges absolutely in an appropriate sense. By a deep result of [Hua95, Hua05], $\Phi_{z,\zeta}$ and $\Psi_{z,\zeta}$ have the same image.

We now assume

$$0 < |z - \zeta| < |\zeta| < |z|, \quad \arg(z - \zeta) = \arg \zeta = \arg z. \quad (1.11)$$

In particular, ζ, z are on the same ray starting from the origin. If we also choose bases $\Xi_{s,k}^l$ and $\Xi_{i,j}^s$ of $\text{Hom}_V(W_s \boxtimes W_k, W_l)$ and $\text{Hom}_V(W_i \boxtimes W_j, W_s)$ respectively, then we have a matrix $\{F_{\gamma\delta}^{\alpha\beta}\}$ (the fusion matrix) representing the invertible map $\Psi_{z,\zeta} \Phi_{z,\zeta}^{-1}$. Equivalently, we have a unique number $F_{\gamma\delta}^{\alpha\beta}$ for each $\alpha, \beta, \gamma, \delta$ such that for each $s \in \mathcal{E}$ and each $\gamma \in \Xi_{s,k}^l, \delta \in \Xi_{i,j}^s$, the fusion relation

$$\mathcal{Y}_\gamma(\mathcal{Y}_\delta(w^{(i)}, z - \zeta) w^{(j)}, \zeta) = \sum_{r \in \mathcal{E}} \sum_{\alpha \in \Xi_{i,r}^l, \beta \in \Xi_{j,k}^r} F_{\gamma\delta}^{\alpha\beta} \cdot \mathcal{Y}_\alpha(w^{(i)}, z) \mathcal{Y}_\beta(w^{(j)}, \zeta) \quad (1.12)$$

holds for each $w^{(i)} \in W_i, w^{(j)} \in W_j$. This fusion matrix is independent of the particular choice of z, ζ satisfying the above mentioned conditions. The associativity isomorphisms of $\text{Rep}(V)$ are defined in such a way that after making $\text{Rep}(V)$ strict, we have

$$\gamma(\delta \otimes \mathbf{1}_k) = \sum_{r \in \mathcal{E}} \sum_{\alpha \in \Xi_{i,r}^l, \beta \in \Xi_{j,k}^r} F_{\gamma\delta}^{\alpha\beta} \cdot \alpha(\mathbf{1}_i \otimes \beta), \quad (1.13)$$

namely, F is also an F -matrix of $\text{Rep}(V)$.

2 Fusion of annihilation and vertex operators

Let W_i, W_j be V -modules. For each $W_s \in \mathcal{E}$, there is a non-degenerate bilinear form $\langle \cdot, \cdot \rangle$ on $\text{Hom}_V(W_i \boxtimes W_j, W_s) \otimes \text{Hom}_V(W_s, W_i \boxtimes W_j)$ such that if $\alpha \in \text{Hom}_V(W_i \boxtimes W_j, W_s)$ and $T \in \text{Hom}_V(W_s, W_i \boxtimes W_j)$, then

$$\alpha T = \langle \alpha, T \rangle \mathbf{1}_s. \quad (2.1)$$

This bilinear form gives an isomorphism

$$\text{Hom}_V(W_s, W_i \boxtimes W_j) \xrightarrow{\cong} \text{Hom}_V(W_i \boxtimes W_j, W_s)^*. \quad (2.2)$$

We shall always identify $\text{Hom}_V(W_s, W_i \boxtimes W_j)$ and $\text{Hom}_V(W_i \boxtimes W_j, W_s)^*$ using the above isomorphism.

Recall from the last section that $\Xi_{i,j}^s$ is a basis of $\text{Hom}_V(W_i \boxtimes W_j, W_s)$. Then we can choose a dual basis $\{\check{\alpha} : \alpha \in \Xi_{i,j}^s\}$. Namely, for each $\alpha \in \Xi_{i,j}^s$, we have $\check{\alpha} \in \text{Hom}_V(W_s, W_i \boxtimes W_j)$, and if $\beta \in \Xi_{i,j}^s$, then $\langle \alpha, \check{\beta} \rangle = \delta_{\alpha,\beta}$. So we also have

$$\alpha \check{\beta} = \delta_{\alpha,\beta} \mathbf{1}_s. \quad (2.3)$$

This implies that

$$\mathbf{1}_{i \boxtimes j} = \sum_{s \in \mathcal{E}} \sum_{\alpha \in \Xi_{i,j}^s} \check{\alpha} \alpha, \quad (2.4)$$

since, by (2.3), the left multiplications of both sides of (2.4) by any $\beta \in \Xi_{i,j}^s$ equal β .

In [HK07], Huang-Kong used the rigidity of $\text{Rep}(V)$ to define a natural isomorphism $\text{Hom}_V(W_{\bar{i}} \boxtimes W_{\bar{j}}, W_{\bar{s}}) \xrightarrow{\cong} \text{Hom}_V(W_i \boxtimes W_j, W_s)^*$. Since $\mathcal{V}(\begin{smallmatrix} \bar{s} \\ \bar{i} \bar{j} \end{smallmatrix})$ is isomorphic to $\mathcal{V}(\begin{smallmatrix} j \\ \bar{i} s \end{smallmatrix})$ by sending \mathcal{Y} to $\mathbb{C}\mathcal{Y}$, we also have an isomorphism

$$\mathfrak{q} : \text{Hom}_V(W_i \boxtimes W_j, W_s)^* \xrightarrow{\cong} \text{Hom}_V(W_{\bar{i}} \boxtimes W_{\bar{j}}, W_{\bar{s}}). \quad (2.5)$$

In the following, we review the construction of this isomorphism.

In [Hua08], Huang showed that $\text{Rep}(V)$ is rigid, and the (categorical) dual object of any V -module W_i could be chosen to be the contragredient module $W_{\bar{i}}$. Moreover, if we define

$$\text{ev}_{\bar{i},i} \in \text{Hom}_V(W_{\bar{i}} \boxtimes W_i, V)$$

such that

$$\mathcal{Y}_{\text{ev}_{\bar{i},i}} = \mathbb{C}\mathcal{Y}_{\kappa(\bar{i})},$$

(Recall that $\mathcal{Y}_{\kappa(\bar{i})}$ is the creation operator of $W_{\bar{i}}$, which is of type $\begin{pmatrix} \bar{i} \\ \bar{i} 0 \end{pmatrix}$. $\mathcal{Y}_{\text{ev}_{\bar{i},i}}$, which is of type $\begin{pmatrix} 0 \\ \bar{i} i \end{pmatrix}$, is called the **annihilation operator** of W_i .) then there is a (unique) morphism

$$\text{coev}_{i,\bar{i}} \in \text{Hom}_V(V, W_i \boxtimes W_{\bar{i}})$$

satisfying the conjugate equations

$$\begin{aligned}(\mathbf{1}_i \otimes \text{ev}_{\bar{i},i})(\text{coev}_{i,\bar{i}} \otimes \mathbf{1}_i) &= \mathbf{1}_i, \\ (\text{ev}_{\bar{i},i} \otimes \mathbf{1}_{\bar{i}})(\mathbf{1}_{\bar{i}} \otimes \text{coev}_{i,\bar{i}}) &= \mathbf{1}_{\bar{i}}.\end{aligned}$$

This is also true for $W_{\bar{i}}$. Thus we have $\text{ev}_{i,\bar{i}}$ and $\text{coev}_{\bar{i},i}$ defined by $\text{ev}_{i,\bar{i}} = \text{ev}_{\bar{i},i}$ and $\text{coev}_{\bar{i},i} = \text{coev}_{i,\bar{i}}$.

Recall the identification (2.2). We define

$$\begin{aligned}\mathfrak{q} : \text{Hom}_V(W_s, W_i \boxtimes W_j) &\xrightarrow{\cong} \text{Hom}_V(W_{\bar{i}} \boxtimes W_s, W_j), \\ T &\mapsto \mathfrak{q}(T) = (\text{ev}_{\bar{i},i} \otimes \mathbf{1}_j)(\mathbf{1}_{\bar{i}} \otimes T).\end{aligned}\tag{2.6}$$

That \mathfrak{q} is an isomorphism follows from the conjugate equations. Using the definition of \mathfrak{q} and equation (2.4), it is easy to see

$$\text{ev}_{\bar{i},i} \otimes \mathbf{1}_j = \sum_{s \in \mathcal{E}} \sum_{\alpha \in \Xi_{i,j}^s} \mathfrak{q}(\check{\alpha})(\mathbf{1}_{\bar{i}} \otimes \alpha).\tag{2.7}$$

Thus, by (1.12) and (1.13), we have the following fusion relation which will play an important role in later sections: Let $z, \zeta \in \mathbb{C}$ satisfy (1.11). Then for any $w^{(i)} \in W_i, w^{(\bar{i})} \in W_{\bar{i}}$,

$$Y_j(\mathcal{Y}_{\text{ev}_{\bar{i},i}}(w^{(\bar{i})}, z - \zeta)w^{(i)}, \zeta) = \sum_{s \in \mathcal{E}} \sum_{\alpha \in \Xi_{i,j}^s} \mathcal{Y}_{\mathfrak{q}(\check{\alpha})}(w^{(\bar{i})}, z)\mathcal{Y}_{\alpha}(w^{(i)}, \zeta).\tag{2.8}$$

Note that for each $s \in \mathcal{E}$, $\{\mathfrak{q}(\check{\alpha}) : \alpha \in \Xi_{i,j}^s\}$ is a basis of $\text{Hom}_V(W_{\bar{i}} \boxtimes W_s, W_j)$. Roughly speaking, this fusion relation says that any intertwining operator arises from fusing the annihilation operators and the vertex operators. This is parallel to the fact that any V -module character occurs in the sum resulting from the modular transformation $\tau \mapsto -1/\tau$ of the vacuum module character.

3 Compressions of intertwining operators

Assume that V is a vertex operator subalgebra (sub-VOA for short) of another CFT-type VOA U with vertex operator Y^U and conformal vector ω . This means that V is a subspace of U , V and U share the same vacuum vector $\mathbf{1}$, and that $Y^U(v_1, z)v_2 = Y(v_1, z)v_2$ when $v_1, v_2 \in V$. Let $L_n^U = Y^U(\omega)_{n+1}$. We shall always assume the additional condition that

$$L_0^U \nu = 2\nu, \quad L_1^U \nu = 0.\tag{3.1}$$

Then by [FZ92] or [LL12] theorem 3.11.12, $(V^c, Y', \mathbf{1}, \nu')$ is a sub-VOA of U , where V^c is the set of all $u \in U$ such that $Y(v)_n u = 0$ for all $v \in V$ and $n \in \mathbb{N}$, Y' is the restriction of Y^U to V^c , and $\nu' = \omega - \nu$. We set $L'_n = Y'(\nu')_{n+1}$.

Assume that V^c is self-dual, CFT-type, and regular. Then $V \otimes V^c$ is also CFT-type and self-dual (and also regular). Thus it is simple. Therefore, the homomorphism of $V \otimes V^c$ -modules

$$V \otimes V^c \rightarrow U, \quad v \otimes v' \mapsto Y(v)_{-1}Y(v')_{-1}\mathbf{1}\tag{3.2}$$

(cf. [LL12] proposition 3.12.7) must be injective. Thus, we can regard $V \otimes V^c$ as a *conformal* sub-VOA of U sharing the same conformal vector $\omega = \nu + \nu' = \nu \otimes \mathbf{1} + \mathbf{1} \otimes \nu'$. Note that by the identification $v \otimes v' = Y(v)_{-1}Y(v')_{-1}\mathbf{1}$, we have $\mathbf{1} = \mathbf{1} \otimes \mathbf{1}$, $v = v \otimes \mathbf{1}$, $v' = \mathbf{1} \otimes v'$.

Recall that by [FHL93] chapter 4, any irreducible $V \otimes V^c$ -module is the tensor product of a V -module and a V^c -module. Moreover, by [ADL05] theorem 2.10, any irreducible intertwining operator of $V \otimes V^c$ can be written as a sum of tensor products of irreducible intertwining operators of V and of V^c . Therefore, any U -module, considered as a $V \otimes V^c$ -module, is a direct sum of those of the form $W_i \otimes W_{i'}$, where W_i is an irreducible V -module and $W_{i'}$ is an irreducible V^c -module. Theorem 2.10 also implies that any intertwining operator of U can be decomposed as a sum of $\mathcal{Y}_\alpha \otimes \mathcal{Y}_{\alpha'}$, where \mathcal{Y}_α and $\mathcal{Y}_{\alpha'}$ are irreducible intertwining operators of V and V^c respectively.

In the following, W_I, W_J, W_K, \dots will denote U -modules, and $W_{i'}, W_{j'}, W_{k'}, \dots$ will denote V^c -modules. $\mathcal{V}^U \binom{K}{I \ J}$ and $\mathcal{V}' \binom{k'}{i' \ j'}$ will denote the corresponding vector spaces of intertwining operators of U and V^c respectively. Note that W_I can not be regarded as a V -module (unless when $\omega = \nu$) but only as a weak V -module (see [DLM97] for the definition.)

Definition 3.1. Let W_i be an irreducible V -module and W_I be a U -module. Let $\varphi : W_i \rightarrow W_I$ and $\psi : W_I \rightarrow W_i$ be homomorphisms of weak V -modules, i.e., they intertwine the actions of V . We say that φ is **grading-preserving** if φ maps each L_0 -eigenspace of W_i into an L_0^U -eigenspace of W_I . We say that ψ is **grading-preserving** if the preimage under ψ of any L_0^U -eigenspace of W_I is contained in an L_0 -eigenspace of W_i .

Remark 3.2. We have seen that there is an identification of $V \otimes V^c$ -modules:

$$W_I \simeq \bigoplus_{s \in \mathcal{E}} W_s \otimes W_{\sigma(s)} \quad (3.3)$$

where for each $s \in \mathcal{E}$, $W_{\sigma(s)}$ is a (non-necessarily irreducible) V^c -module. We can also regard (3.3) as a decomposition of W_I into irreducible weak V -modules, where for each $s \in \mathcal{E}$, $W_{\sigma(s)}$ is the multiplicity space of W_s . (Note that different elements in \mathcal{E} give rise to non-equivalent irreducible modules.) Choose any $s \in \mathcal{E}$. Choose $\varphi : W_s \rightarrow W_I$ and $\psi : W_I \rightarrow W_s$ to be homomorphisms of weak V -modules. Then it is not hard to see that we can find $w^{(\sigma(s))} \in W_{\sigma(s)}$ and $\varpi \in W_{\sigma(s)}^*$ (note that $W_{\sigma(s)}^*$ is the dual vector space of $W_{\sigma(s)}$) such that

$$\varphi = \mathbf{1}_s \otimes w^{(\sigma(s))}, \quad \psi = \mathbf{1}_s \otimes \varpi, \quad (3.4)$$

where $w^{(\sigma(s))}$ is considered as the linear map $\mathbb{C} \rightarrow W_{\sigma(s)}$ sending 1 to $w^{(\sigma(s))}$.² Moreover, outside $W_s \otimes W_{\sigma(s)}$, $\mathbf{1}_s \otimes \varpi$ is defined to be the zero functional. Thus, it is clear that φ (resp. ψ) is *grading-preserving* if and only if $w^{(\sigma(s))}$ (resp. ϖ) equals an (L_0') homogeneous vector of $W_{\sigma(s)}$ (resp. $W_{\sigma(s)}^*$).

²To see that φ can be written in this way, choose any $t \in \mathcal{E}$ and $\omega \in W_{\sigma(t)}^*$, and consider the homomorphism of irreducible V -modules $T_\omega : W_s \rightarrow W_t$ defined by $T_\omega = (\mathbf{1}_t \otimes \omega) \circ \varphi$. Then $T_\omega = 0$ whenever $s \neq t$ (since $W_s \neq W_t$). So the image of φ is in $W_s \otimes W_{\sigma(s)}$. Now assume $t = s$. Then T_ω is a scalar. Choose a basis $\{e_1, e_2, \dots\}$ of W_s , and write $\varphi(e_1) = \sum_n e_n \otimes w_n$ where each w_n is in $W_{\sigma(s)}$. Then, for each ω , $T_\omega(e_1) = \sum_n \omega(w_n)e_n$ is a scalar multiple of e_1 , which shows that $w_n = 0$ when $n > 1$. Thus $\varphi = \mathbf{1}_s \otimes w_1$. That ψ has the desired form can be proved similarly.

Definition 3.3. Let W_i be an irreducible V -module and W_I be a U -module. We say that W_i is a **compression** of W_I if W_I (considered as a weak V -module) has an irreducible weak V -submodule isomorphic to W_i . Equivalently, the $V \otimes V^c$ -module W_I has a (non-trivial) irreducible submodule isomorphic to $W_i \otimes W_{i'}$ for some irreducible V^c -module $W_{i'}$.

Definition 3.4. Let $\mathcal{Y} \in \mathcal{V} \binom{k}{i \ j}$ be an irreducible intertwining operator of V .

(a) Let $\mathcal{Y}^U \in \mathcal{V}^U \binom{K}{I \ J}$ be an intertwining operator of U . We say that \mathcal{Y} is a **compression** of \mathcal{Y}^U , if there exist $\lambda \in \mathbb{Q}$ and grading-preserving homomorphisms of weak V -modules $\varphi : W_i \rightarrow W_I$, $\phi : W_j \rightarrow W_J$, and $\psi : W_K \rightarrow W_{k'}$, such that for any $w^{(i)} \in W_i$ and $z \in \mathbb{C}^\times$

$$\mathcal{Y}(w^{(i)}, z) = z^\lambda \cdot \psi \mathcal{Y}^U(\varphi w^{(i)}, z) \phi.$$

(b) If W_I, W_J are U -modules, we say that \mathcal{Y} is a **compression of type $\binom{\bullet}{I \ J}$ intertwining operators of U** , if \mathcal{Y} is a (finite) sum of compressions of intertwining operators of U whose charge spaces are W_I and source spaces are W_J .

(c) If \mathcal{Y} is a (finite) sum of compressions of intertwining operators of U , we simply say that \mathcal{Y} is a **compression of intertwining operators of U**

Proposition 3.5. Let W_I, W_J, W_K be U -modules with $V \otimes V^c$ -irreducible decompositions

$$W_I \simeq \bigoplus W_i \otimes W_{i'}, \quad W_J \simeq \bigoplus W_j \otimes W_{j'}, \quad W_K \simeq \bigoplus W_k \otimes W_{k'}.$$

Then, according to these decompositions, any $\mathcal{Y}^U \in \mathcal{V}^U \binom{K}{I \ J}$ can be written as a sum of elements of the form $\mathcal{Y} \otimes \mathcal{Y}'$, where $\mathcal{Y} \in \mathcal{V} \binom{k}{i \ j}$ is the compression of a type $\binom{K}{I \ J}$ intertwining operator of U , and \mathcal{Y}' is an irreducible intertwining operator of V^c .

Proof. We fix irreducible $V \otimes V^c$ -submodules $W_i \otimes W_{i'}, W_j \otimes W_{j'}, W_k \otimes W_{k'}$ of W_I, W_J, W_K respectively. Let $\Theta_{i', j'}^{k'}$ be a basis of $\text{Hom}_{V^c}(W_{i'} \boxtimes W_{j'}, W_{k'})$. We have a (functorial) isomorphism

$$\text{Hom}_{V^c}(W_{i'} \boxtimes W_{j'}, W_{k'}) \xrightarrow{\simeq} \mathcal{V}' \binom{k'}{i' \ j'}, \quad \alpha' \mapsto \mathcal{Y}'_{\alpha'} \quad (3.5)$$

similar to (1.2). Consider \mathcal{Y}^U as an intertwining operator of $V \otimes V^c$, and restrict it to $W_i \otimes W_{i'}, W_j \otimes W_{j'}, W_k \otimes W_{k'}$. Then, by [ADL05] theorem 2.10, this restriction is a sum of tensor products of V - and V^c -intertwining operators. Assume without loss of generality that this restriction is non-zero. Then $W_k, W_{k'}$ must be irreducible submodules of $W_i \boxtimes W_j, W_{i'} \boxtimes W_{j'}$ respectively. Now, for each $\alpha' \in \Theta_{i', j'}^{k'}$, we can find $\alpha \in \text{Hom}_V(W_i \boxtimes W_j, W_k)$ (not necessarily in $\Xi_{i, j}^k$) such that the restriction of \mathcal{Y}^U equals

$$\sum_{\alpha' \in \Theta_{i', j'}^{k'}} \mathcal{Y}_\alpha \otimes \mathcal{Y}'_{\alpha'}.$$

We shall show that each \mathcal{Y}_α is a sum of compressions of type $\binom{K}{I \ J}$ intertwining operators of U .

Choose $\alpha' \in \Theta_{i',j'}^{k'}$, and apply corollary 1.2 to V^c . Then there exist homogeneous vectors $w_1^{(i')}, \dots, w_m^{(i')} \in W_{i'}, w_1^{(j')}, \dots, w_m^{(j')} \in W_{j'}, w^{(\bar{k}')} \in W_{\bar{k}'}$, and constants $\lambda_1, \dots, \lambda_m \in \mathbb{Q}$, such that for any $\beta' \in \Theta_{i',j'}^{k'}$, the expression

$$\sum_{l=1}^m z^{\lambda_l} \langle \mathcal{Y}'_{\beta'}(w_l^{(i')}, z) w_l^{(j')}, w^{(\bar{k}')} \rangle \quad (3.6)$$

is a constant (over the complex variable z), and this constant is non-zero if and only if $\beta' = \alpha'$. By scaling the vector $w^{(\bar{k}')}$, we may assume that when $\beta' = \alpha'$, the above constant is 1. Now, for each $l = 1, 2, \dots, m$, define grading-preserving homomorphisms of weak V -modules $\varphi_l : W_i \rightarrow W_I, \phi_l : W_j \rightarrow W_J, \psi : W_K \rightarrow W_k$ by

$$\varphi_l = \mathbf{1}_i \otimes w_l^{(i')}, \quad \phi_l = \mathbf{1}_j \otimes w_l^{(j')}, \quad \psi = \mathbf{1}_k \otimes w^{(\bar{k}')}.$$

Then we have

$$\mathcal{Y}_\alpha(w^{(i)}, z) = \sum_{l=1}^m z^{\lambda_l} \cdot \psi \mathcal{Y}^U(\varphi_l w^{(i)}, z) \phi_l.$$

This finishes the proof. \square

4 Proof of the main result

In this section, we assume that $V^{cc} = V$. Let W_I, W_J be V -modules, and we fix irreducible decompositions of $V \otimes V^c$ -modules:

$$U \simeq (V \otimes V^c) \oplus \left(\bigoplus W_a \otimes W_{a'} \right), \quad (4.1)$$

$$W_I \simeq (V \otimes V^c) \oplus \left(\bigoplus W_i \otimes W_{i'} \right), \quad (4.2)$$

$$W_J \simeq (V \otimes V^c) \oplus \left(\bigoplus W_j \otimes W_{j'} \right). \quad (4.3)$$

We first recall the following obvious fact.

Proposition 4.1. *If $W_a \otimes W_{a'}$ is an irreducible $V \otimes V^c$ -submodule of $U \ominus (V \otimes V^c)$, then W_a is not isomorphic to V and $W_{a'}$ is not isomorphic to V^c .*

Proof. If $W_{a'}$ is isomorphic to V^c , then $W_{a'}$ contains a non-zero homogeneous vector w_2 equivalent to the vacuum vector of V^c . Choose any non-zero $w_1 \in W_a$. Then for any $v' \in V^c$ and $n \in \mathbb{N}$, $Y'(v')_n w_2 = 0$. Therefore $Y^U(v')_n(w_1 \otimes w_2) = w_1 \otimes Y'(v')_n w_2 = 0$. Thus $w_1 \otimes w_2 \in V^{cc} = V = V \otimes \mathbf{1}$, which is impossible since $w_1 \otimes w_2$ is not in $V \otimes V^c$. So $W_{a'}$ is not isomorphic to V^c . Since $V^{ccc} = V^c$, for a similar reason, W_a is also not isomorphic to V . \square

For each irreducible V^c -module, we choose a representing element, and let them form a finite set \mathcal{E}' . Assume $W_{0'} := V^c$ is in \mathcal{E}' . If $W_{i'}, W_{j'}, W_{k'}$ are V^c -modules, we choose a basis $\Theta_{i,j}^k$ of $\text{Hom}_{V^c}(W_{i'} \boxtimes W_{j'}, W_{k'})$. The linear isomorphism

$$\eta : \text{Hom}_{V^c}(W_{i'} \boxtimes W_{j'}, W_{k'})^* \xrightarrow{\cong} \text{Hom}_{V^c}(W_{\bar{i}'} \boxtimes W_{\bar{j}'}, W_{\bar{k}'}) \quad (4.4)$$

and the morphism

$$\text{ev}_{\bar{i},i'} \in \text{Hom}_{V^c}(W_{\bar{i}} \boxtimes W_{i'}, V^c)$$

are defined as in section 2. Then $\mathcal{Y}'_{\text{ev}_{\bar{i},i'}}$ is the annihilation operator of $W_{i'}$.

According to the decomposition for W_I , $W_{\bar{I}}$ also has the corresponding decomposition:

$$W_{\bar{I}} \simeq (V \otimes V^c) \oplus \left(\bigoplus W_{\bar{i}} \otimes W_{i'} \right).$$

Let

$$\text{ev}_{\bar{I},I} \in \text{Hom}_U(W_{\bar{I}} \boxtimes W_I, U)$$

which corresponds to the annihilation operator $\mathcal{Y}_{\text{ev}_{\bar{I},I}}^U$ of the U -module W_I . Suppose that in the above decompositions, $W_{\bar{i}} \boxtimes W_{i'}$ is an irreducible submodule of $W_I \ominus (V \otimes V^c)$. If we regard $\mathcal{Y}_{\text{ev}_{\bar{I},I}}^U$ as an intertwining operator of $V \otimes V^c$, then it is easy to see that the restriction of $\mathcal{Y}_{\text{ev}_{\bar{I},I}}^U$ to the charge subspace $W_{\bar{i}} \boxtimes W_{i'}$ source subspace $W_{\bar{i}} \boxtimes W_{i'}$ and the target subspace $V \otimes V^c$ is $\mathcal{Y}_{\text{ev}_{\bar{i},i}} \otimes \mathcal{Y}'_{\text{ev}_{i',i'}}$, the tensor product of the annihilation operators of $W_{\bar{i}}$ and of $W_{i'}$.

Theorem 4.2. *Let V be a vertex operator subalgebra of U satisfying (3.1). Assume that U , V , and V^c are CFT type, self-dual, and regular VOAs. Assume also that $V^{cc} = V$. Let W_I, W_J be U -modules. Let W_i, W_j be irreducible V -modules that are compressions of W_I and W_J respectively. Then any irreducible intertwining operator of V with charge space W_i and source space W_j is a compression of type $\binom{\bullet}{I \ J}$ intertwining operators of U .*

Proof. Let $W_i \otimes W_{i'}$ and $W_j \otimes W_{j'}$ be irreducible $V \otimes V^c$ -submodules of W_I, W_J respectively. Assume that $k \in \mathcal{E}$ and not all type $\binom{k}{i \ j}$ intertwining operators of V are compressions of type $\binom{\bullet}{I \ J}$ intertwining operators of U . Let $\mathcal{V} \binom{k}{i \ j}$ be a subspace of $\mathcal{V} \binom{k}{i \ j}$ with codimension 1 containing all elements of $\mathcal{V} \binom{k}{i \ j}$ that are compressions of type $\binom{\bullet}{I \ J}$ intertwining operators of U . Choose a nonzero element $\mathfrak{A} \in \text{Hom}_V(W_i \boxtimes W_j, W_k)$ such that $\mathfrak{A} \notin \mathcal{V} \binom{k}{i \ j}$. We assume that the basis $\Xi_{i,j}^k$ of $\text{Hom}_V(W_i \boxtimes W_j, W_k)$ is chosen such that $\mathfrak{A} \in \Xi_{i,j}^k$, and that $\mathcal{Y}_\alpha \in \mathcal{V} \binom{k}{i \ j}$ for any $\alpha \in \Xi_{i,j}^k$ not equal to \mathfrak{A} .

Choose $z, \zeta \in \mathbb{C}$ satisfying (1.11). Recall that Y_I^U is the U -vertex operator of W_I and $\mathcal{Y}_{\text{ev}_{\bar{I},I}}^U$ is the U -annihilation operator of W_I . In the following, we shall calculate the fusion relation for the iterate of $V \otimes V^c$ -intertwining operators Y_J^U and $\mathcal{Y}_{\text{ev}_{\bar{I},I}}^U$ (with restricted charge, source, and target spaces) in two ways. These two methods will give incompatible results, which therefore lead to a contradiction. Let $\pi_{j \otimes j'}$ be the projection of the algebraic completion of W_J onto the one of $W_j \otimes W_{j'}$.

Step 1. Note that for each $s \in \mathcal{E}$, $s' \in \mathcal{E}'$, the set $\Xi_{i,j}^s \times \Theta_{i',j'}^{s'}$ (more precisely, $\{\mathcal{Y}_\alpha \otimes \mathcal{Y}'_{\alpha'} : \alpha \in \Xi_{i,j}^s, \alpha' \in \Theta_{i',j'}^{s'}\}$) is a basis of the vector space of type $\binom{W_s \otimes W_{s'}}{W_i \otimes W_{i'} \ W_j \otimes W_{j'}}$ intertwining operators of $V \otimes V^c$. (See [ADL05] theorem 2.10; it is also an easy consequence of corollary 1.2.) Thus, for any $s \in \mathcal{E}$, $s' \in \mathcal{E}'$ and $\alpha, \beta \in \text{Hom}_V(W_i \boxtimes W_j, W_s)$, $\alpha', \beta' \in \text{Hom}_{V^c}(W_{i'} \boxtimes W_{j'}, W_{s'})$, there is a unique constant $\lambda_{\alpha,\beta,\alpha',\beta'} \in \mathbb{C}$ such that for any

$$w_1 \in W_{\bar{i}}, w_2 \in W_{\bar{i}'}, w_3 \in W_i, w_4 \in W_{i'}, w_5 \in W_j, w_6 \in W_{j'}, \quad (4.5)$$

the following fusion relation of $V \otimes V^c$ -intertwining operators holds:

$$\begin{aligned} & \pi_{j \otimes j'} \cdot Y_J^U (\mathcal{Y}_{\text{ev}_{\bar{T}, I}}^U (w_1 \otimes w_2, z - \zeta)(w_3 \otimes w_4), \zeta)(w_5 \otimes w_6) \\ = & \sum_{\substack{s \in \mathcal{E} \\ s' \in \mathcal{E}'}} \sum_{\substack{\alpha, \beta \in \Xi_{i, j}^s \\ \alpha', \beta' \in \Theta_{i', j'}^{s'}}} \lambda_{\alpha, \beta, \alpha', \beta'} \cdot \mathcal{Y}_{\mathfrak{q}(\check{\beta})}(w_1, z) \mathcal{Y}_{\alpha}(w_3, \zeta) w_5 \otimes \mathcal{Y}'_{\mathfrak{q}(\check{\beta}')} (w_2, z) \mathcal{Y}'_{\alpha'}(w_4, \zeta) w_6. \end{aligned} \quad (4.6)$$

(Recall (1.2) and (3.5) for the notations $\mathcal{Y}, \mathcal{Y}'$.) On the other hand, by (2.8), the iterate of the U -intertwining operators Y_J^U and $\mathcal{Y}_{\text{ev}_{\bar{T}, I}}^U$ equals a sum of products of type $\binom{J}{\bar{T} \bullet}$ intertwining operators and type $\binom{\bullet}{I J}$ intertwining operators of U . Therefore, by proposition 3.5 and the uniqueness of fusion coefficients, we have

$$\lambda_{\mathfrak{q}, \beta, \alpha', \beta'} = 0 \quad (4.7)$$

for any $s' \in \mathcal{E}'$, $\beta \in \Xi_{i, j'}^k$, and $\alpha', \beta' \in \Theta_{i', j'}^{s'}$. In particular, the right hand side of (4.6) has no terms containing $\mathcal{Y}_{\mathfrak{q}(\check{\alpha})}(w_1, z) \mathcal{Y}_{\mathfrak{q}}(w_3, \zeta) w_5$.

Step 2. We calculate the iterate of $\pi_{j \otimes j'} \cdot Y_J^U$ and $\mathcal{Y}_{\text{ev}_{\bar{T}, I}}^U$ using a different method, and show that some terms containing $\mathcal{Y}_{\mathfrak{q}(\check{\alpha})}(w_1, z) \mathcal{Y}_{\mathfrak{q}}(w_3, \zeta) w_5$ will appear. By the paragraph before the theorem, we know that for any w_1, w_2, \dots, w_6 as in (4.5),

$$\begin{aligned} & \mathcal{Y}_{\text{ev}_{\bar{T}, I}}^U (w_1 \otimes w_2, z - \zeta)(w_3 \otimes w_4) \\ = & \mathcal{Y}_{\text{ev}_{\bar{i}, i}}(w_1, z - \zeta) w_3 \otimes \mathcal{Y}'_{\text{ev}_{\bar{i}', i'}}(w_2, z - \zeta) w_4 \\ & + \sum_{W_a \otimes W_{a'}} \sum_{\gamma, \gamma'} \mathcal{Y}_{\gamma}(w_1, z - \zeta) w_3 \otimes \mathcal{Y}'_{\gamma'}(w_2, z - \zeta) w_4 \end{aligned} \quad (4.8)$$

where the first sum is over all irreducible $V \otimes V^c$ -submodules of $U \ominus (V \otimes V^c)$ as in the decomposition (4.1), $\gamma \in \text{Hom}_V(W_{\bar{i}} \boxtimes W_i, W_a)$, and $\gamma' \in \text{Hom}_{V^c}(W_{\bar{i}'} \boxtimes W_{i'}, W_{a'})$. We shall now calculate the iterate of $\pi_{j \otimes j'} \cdot Y_J^U$ with each term on the right hand side of (4.8).

The first term is in the algebraic completion of $V \otimes V^c$. Moreover, the restriction of Y_J^U (regarded as a $V \otimes V^c$ -intertwining operator) to $V \otimes V^c, W_j \otimes W_{j'}, W_j \otimes W_{j'}$ equals $Y_j \otimes Y_{j'}$, where $Y_j, Y_{j'}$ are respectively the vertex operators of the V -module W_j and the V^c -module $W_{j'}$. Therefore, by (2.8),

$$\begin{aligned} & \pi_{j \otimes j'} \cdot Y_J^U (\mathcal{Y}_{\text{ev}_{\bar{i}, i}}(w_1, z - \zeta) w_3 \otimes \mathcal{Y}'_{\text{ev}_{\bar{i}', i'}}(w_2, z - \zeta) w_4, \zeta)(w_5 \otimes w_6) \\ = & Y_j (\mathcal{Y}_{\text{ev}_{\bar{i}, i}}(w_1, z - \zeta) w_3, \zeta) w_5 \otimes Y_{j'} (\mathcal{Y}'_{\text{ev}_{\bar{i}', i'}}(w_2, z - \zeta) w_4, \zeta) w_6 \\ = & \sum_{s \in \mathcal{E}} \sum_{\alpha \in \Xi_{i, j}^s} \mathcal{Y}_{\mathfrak{q}(\check{\alpha})}(w_1, z) \mathcal{Y}_{\alpha}(w_3, \zeta) w_5 \otimes Y_{j'} (\mathcal{Y}'_{\text{ev}_{\bar{i}', i'}}(w_2, z - \zeta) w_4, \zeta) w_6. \end{aligned} \quad (4.9)$$

In the above expression, (the sum of) all the terms containing $\mathcal{Y}_{\mathfrak{q}(\check{\alpha})}(w_1, z) \mathcal{Y}_{\mathfrak{q}}(w_3, \zeta) w_5$ is

$$\mathcal{Y}_{\mathfrak{q}(\check{\alpha})}(w_1, z) \mathcal{Y}_{\mathfrak{q}}(w_3, \zeta) w_5 \otimes Y_{j'} (\mathcal{Y}'_{\text{ev}_{\bar{i}', i'}}(w_2, z - \zeta) w_4, \zeta) w_6. \quad (4.10)$$

On the other hand, suppose that when restricted to the charge subspace $W_a \otimes W_{a'}$ (where $W_a \otimes W_{a'}$ is an irreducible submodule of $U \ominus (V \otimes V^c)$) and source and target

subspace $W_j \otimes W_{j'}$, the $V \otimes V^c$ -intertwining operator Y_J^U could be written as $\sum_{\delta, \delta'} \mathcal{Y}_\delta \otimes \mathcal{Y}'_{\delta'}$, where each \mathcal{Y}_δ is of type $\binom{j}{a \ j}$ and $\mathcal{Y}'_{\delta'}$ has type $\binom{j'}{a' \ j'}$. Then the iterate of $\pi_{j \otimes j'} \cdot Y_J^U$ with the second term of (4.8) is

$$\begin{aligned} & \sum_{W_a \otimes W_{a'}} \sum_{\gamma, \gamma'} \pi_{j \otimes j'} \cdot Y_J^U (\mathcal{Y}_\gamma(w_1, z - \zeta)w_3 \otimes \mathcal{Y}_{\gamma'}(w_2, z - \zeta)w_4, \zeta)(w_5 \otimes w_6) \\ &= \sum_{W_a \otimes W_{a'}} \sum_{\substack{\gamma, \gamma' \\ \delta, \delta'}} \mathcal{Y}_\delta (\mathcal{Y}_\gamma(w_1, z - \zeta)w_3, \zeta)w_5 \otimes \mathcal{Y}'_{\delta'} (\mathcal{Y}'_{\gamma'}(w_2, z - \zeta)w_4, \zeta)w_6. \end{aligned} \quad (4.11)$$

If we write each $\mathcal{Y}_\delta (\mathcal{Y}_\gamma(w_1, z - \zeta)w_3, \zeta)w_5$ as a sum of products of V -intertwining operators under the bases $\Xi_{i,j}^s$ and $\{\mathfrak{q}(\check{\alpha}) : \alpha \in \Xi_{i,j}^s\}$ (over all $s \in \mathcal{E}$) similar to part of (4.6), then the sum of all the terms containing $\mathcal{Y}_{\mathfrak{q}(\check{\alpha})}(w_1, z)\mathcal{Y}_{\mathfrak{q}}(w_3, \zeta)w_5$ should be

$$\mathcal{Y}_{\mathfrak{q}(\check{\alpha})}(w_1, z)\mathcal{Y}_{\mathfrak{q}}(w_3, \zeta)w_5 \otimes \sum_{W_a \otimes W_{a'}} \sum_{\substack{\gamma, \gamma' \\ \delta, \delta'}} \kappa_{\gamma, \delta} \cdot \mathcal{Y}'_{\delta'} (\mathcal{Y}'_{\gamma'}(w_2, z - \zeta)w_4, \zeta)w_6 \quad (4.12)$$

where each $\kappa_{\gamma, \delta}$ is a constant. By proposition 4.1, every $W_{a'}$ (which is irreducible) is not isomorphic to V^c . Therefore, as the linear map $\Phi_{z, \zeta}$ (see (1.9)) is injective, the sum of (4.10) and (4.12) is not zero for some w_1, \dots, w_6 satisfying (4.5). This shows that (4.6) (which is the sum of (4.9) and (4.11)) has non-zero terms containing $\mathcal{Y}_{\mathfrak{q}(\check{\alpha})}(w_1, z)\mathcal{Y}_{\mathfrak{q}}(w_3, \zeta)w_5$. In other words, $\lambda_{\mathfrak{q}, \mathfrak{q}, \alpha', \beta'} \neq 0$ for some $s' \in \mathcal{E}'$ and $\alpha', \beta' \in \Theta_{i', j'}^{s'}$. This gives a contradiction. \square

The following result was proved in [KM15]:

Theorem 4.3. *Let V be a vertex operator subalgebra of U satisfying (3.1). Assume that U , V , and V^c are CFT type, self-dual, and regular VOAs. Assume also that $V^{cc} = V$. Then any irreducible V -module is the compression of a U -module.*

The above two theorems imply immediately the following:

Theorem 4.4. *Under the assumption of theorem 4.3, any irreducible intertwining operator of V is a compression of intertwining operators of U .*

References

- [ABD04] Abe, T., Buhl, G. and Dong, C., 2004. Rationality, regularity, and C_2 -cofiniteness. Transactions of the American Mathematical Society, 356(8), pp.3391-3402.
- [ACL19] Arakawa, T., Creutzig, T. and Linshaw, A.R., 2019. W -algebras as coset vertex algebras. Inventiones mathematicae, 218(1), pp.145-195.
- [ADL05] Abe, T., Dong, C., Li, H. 2005. Fusion Rules for the Vertex Operator Algebras $M(1)^+$ and V_L^+ . Comm. Math. Phys, 253, pp.171-219.
- [AM88] Anderson, G. and Moore, G., 1988. Rationality in conformal field theory. Communications in mathematical physics, 117(3), pp.441-450.
- [Ara15a] Arakawa, T., 2015. Associated varieties of modules over Kac–Moody algebras and C_2 -cofiniteness of W -algebras. International Mathematics Research Notices, 2015(22), pp.11605-11666.

- [Ara15b] Arakawa, T., 2015. Rationality of W -algebras: principal nilpotent cases. *Annals of Mathematics*, pp.565-604.
- [CKLW18] Carpi, S., Kawahigashi, Y., Longo, R. and Weiner, M., 2018. From vertex operator algebras to conformal nets and back (Vol. 254, No. 1213). *Memoir of the American Mathematical Society*.
- [DLM97] Dong, C., Li, H. and Mason, G., 1997. Regularity of rational vertex operator algebras. *Advances in Mathematics*, 132(1), pp.148-166.
- [DLM00] Dong, C., Li, H. and Mason, G., 2000. Modular-Invariance of Trace Functions in Orbifold Theory and Generalized Moonshine. *Communications in Mathematical Physics*, 214(1), pp.1-56.
- [FHL93] Frenkel, I., Huang, Y.Z. and Lepowsky, J., 1993. On axiomatic approaches to vertex operator algebras and modules (Vol. 494). *American Mathematical Soc.*
- [FZ92] Frenkel, I.B. and Zhu, Y., 1992. Vertex operator algebras associated to representations of affine and Virasoro algebras. *Duke Mathematical Journal*, 66(1), pp.123-168.
- [Gui18] Gui, B., 2018. Categorical extensions of conformal nets. *arXiv preprint arXiv:1812.04470*.
- [Gui19a] Gui, B., 2019. Unitarity of the modular tensor categories associated to unitary vertex operator algebras, I, *Comm. Math. Phys.*, 366(1), pp.333-396.
- [Gui19b] Gui, B., 2019. Energy bounds condition for intertwining operators of type B , C , and G_2 unitary affine vertex operator algebras. *Trans. Amer. Math. Soc.* 372 (2019), 7371-7424
- [Gui20] Gui, B., 2020. Unbounded field operators in categorical extensions of conformal nets. *arXiv preprint arXiv:2001.03095*.
- [HK07] Huang, Y.Z. and Kong, L., 2007. Full field algebras. *Communications in mathematical physics*, 272(2), pp.345-396.
- [HL13] Huang, Y.Z. and Lepowsky, J., 2013. Tensor categories and the mathematics of rational and logarithmic conformal field theory. *Journal of Physics A: Mathematical and Theoretical*, 46(49), p.494009.
- [Hua95] Huang, Y.Z., 1995. A theory of tensor products for module categories for a vertex operator algebra, IV. *Journal of Pure and Applied Algebra*, 100(1-3), pp.173-216.
- [Hua05] Huang, Y.Z., 2005. Differential equations and intertwining operators. *Communications in Contemporary Mathematics*, 7(03), pp.375-400.
- [Hua08] Huang, Y.Z., 2008. Rigidity and modularity of vertex tensor categories. *Communications in contemporary mathematics*, 10(supp01), pp.871-911.
- [KM15] Krauel, M. and Miyamoto, M., 2015. A modular invariance property of multivariable trace functions for regular vertex operator algebras. *Journal of Algebra*, 444, pp.124-142.
- [LL12] Lepowsky, J. and Li, H., 2012. Introduction to vertex operator algebras and their representations (Vol. 227). *Springer Science & Business Media*.
- [Ten17] Tener, J.E., 2017. Construction of the unitary free fermion Segal CFT. *Communications in Mathematical Physics*, 355(2), pp.463-518.
- [Ten19a] Tener, J.E., 2019. Geometric realization of algebraic conformal field theories. *Advances in Mathematics*, 349, pp.488-563.
- [Ten19b] Tener, J.E., 2019. Representation theory in chiral conformal field theory: from fields to observables. *Selecta Mathematica*, 25(5), p.76.
- [Ten19c] Tener, J.E., 2019. Fusion and positivity in chiral conformal field theory. *arXiv preprint arXiv:1910.08257*.
- [TL04] Toledano-Laredo, V., 2004. Fusion of positive energy representations of $LSpin_{2n}$. *arXiv preprint math/0409044*.
- [Was98] Wassermann, A., 1998. Operator algebras and conformal field theory III. Fusion of positive energy representations of $LSU(N)$ using bounded operators. *Inventiones mathematicae*, 133(3), pp.467-538.

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, USA.

E-mail: bin.gui@rutgers.edu binguimath@gmail.com